



## Reviews

### Results of Lead Research: Prenatal Exposure and Neurological Consequences

Robert A. Goyer

National Institute of Environmental Health Sciences, Research Triangle Park, NC 27709 USA

The history of advances in the understanding of the toxic effects of lead over the past 20 years is an outstanding example of how knowledge learned from research can impact public health. Measures that have had the greatest impact on reducing exposure to lead are reduction of lead from gasoline, elimination of lead solder from canned food, removal of lead from paint, and abatement of housing containing lead-based paint. Nevertheless, continuing factors that enhance risk to lead exposure, particularly during fetal life, are low socioeconomic status, old housing with lead-containing paint, and less than ideal nutrition, particularly low dietary intake of calcium, iron, and zinc. Prenatal exposure may result from endogenous sources such as lead in the maternal skeletal system or maternal exposures from diet and the environment. Experimental studies have shown that the developing nervous system is particularly sensitive to the toxic effects of lead and that a large number of the effects in the nervous system are due to interference of lead with biochemical functions dependent on calcium ions and impairment of neuronal connections dependent on dendritic pruning. There is need for more study to determine whether these effects are a continuum of prenatal lead exposure or whether prenatal exposure to lead produces unique effects. *Key words:* blood lead levels, fetal brain, fetal susceptibility, lead, mechanisms of central nervous system effects, neurological effects, pregnancy, prenatal exposure, risk factors. *Environ Health Perspect* 104:1050–1054 (1996)

Lead toxicity is one of the most studied environmental health issues and is also the most outstanding example of how knowledge learned from research can impact public health. The National Institute of Environmental Health Sciences (NIEHS) has provided support for many of the basic studies over the past 30 years, resulting in advances in understanding cellular effects of lead and effects on organ systems. Many of these accomplishments have been summarized in the NIEHS-sponsored conferences published in *Environmental Health Perspectives* (1) and in other published reports (2–4).

#### Research Accomplishments

The most significant research accomplishments have concerned effects of low-level exposures to lead on cognitive and behavioral development of young children. The 1992 World Health Organization/International Program for Chemical Safety Task Group on Effects of Inorganic Lead concluded that blood lead levels in young children generally below 25  $\mu\text{g}/\text{dl}$  are associated with a reduction in IQ scores (4). The size of the apparent IQ effect as assessed at 4 years of age and above is a deficit between 1 and 5 IQ points for each 10  $\mu\text{g}/\text{dl}$  increment in blood lead,

with a likely size effect between 1 and 3 points. This conclusion is based on a large number of retrospective and prospective epidemiological studies and is similar to the 1991 statement from the U.S. Centers for Disease Control (5) and other national and international groups (6,7). Results of epidemiological studies to date do not provide evidence of a threshold. It is suggested that, with blood lead levels below 10  $\mu\text{g}/\text{dl}$ , the effect of confounding variables and lack of precision of analytical and psychological measurements limits the ability to detect effects below the 10–15  $\mu\text{g}/\text{dl}$  range. Animal studies support a causal relationship between lead and nervous system effects, and there are reports of intellectual deficits in monkeys and rats with blood lead levels in the 11–15  $\mu\text{g}/\text{dl}$  range (8).

The result of the present awareness of lead effects has been to adopt measures that reduce human exposure to lead and reduce risks to related health effects. In 1990 and 1991, the U.S. Departments of Health and Human Services and Housing and Urban Development and the U.S. Environmental Protection Agency (EPA) published strategy documents outlining measures to reduce lead exposure (9–11). The measure with the probably greatest impact was the reduc-

tion of lead in gasoline, although the reasons for this action were based on broader environmental considerations. There has also been a reduction of lead in food due to reduced use of solders in food and soft drink cans. The Food and Drug Administration (FDA) Market Basket Surveys indicate that the typical daily intake of lead for a 2-year-old child dropped from 30  $\mu\text{g}/\text{day}$  in 1982 to about 2  $\mu\text{g}/\text{day}$  in 1991 (12). A federal law provides a Medicaid benefit package for all children through the Early and Periodic Screening Diagnosis and Treatment program for lead-poisoned children, including chelation therapy, vitamin and mineral supplementation, medical care, environmental investigations, education services, and nutritional and developmental assessments (13).


Research achievements that contribute to prevention of disease are not as dramatic as the discovery or implementation of cures for existent disease but are certainly no less important. Research regarding the health effects of lead may make a small contribution that helps millions of people, but is probably unknown to them. While these beneficial effects may not be apparent to the individuals who are spared the problems of toxicity, the implementation of measures to reduce exposure to lead are clearly visible when examined on a population basis.

Figure 1 compares blood lead levels in children 1–5 years of age in the United States as measured by the National Health and Nutrition Examination Survey (NHANES) II in 1976–1980 (Fig. 1A) to blood lead levels as measured in children by NHANES III, 1988–1991 (Fig. 1B) (14). In this 10-year period, the mean blood lead for persons in

Address correspondence to: R. A. Goyer, National Institute of Environmental Health Sciences, PO Box 12233, Research Triangle Park, NC 27709 USA.

Adapted from presentations at the annual meeting of the American Industrial Hygiene Association, Kansas City, 25 May 1995, and the Vth COM-TOX Symposium on Toxicology and Chemistry of Metals, Vancouver, British Columbia, 10–13 July 1995.

Received 11 March 1996; accepted 4 June 1996.



**Figure 1.** Blood lead levels in children 1–5 years of age in the United States as measured by NHANES II in 1976–1980 (A) versus blood lead levels as measured in children by NHANES III, 1988–1991 (B). From Pirkle et al. (15).

the general population of all ages decreased by 78%, from 12.8 to 2.8  $\mu\text{g}/\text{dl}$  between 1976 and 1991, and 77% for children ages 1–5 years, from 15  $\mu\text{g}/\text{dl}$  to 3.6  $\mu\text{g}/\text{dl}$ .

In spite of measures reducing lead exposure to date, large numbers of children in the United States have high exposure to lead and are at risk for impaired cognitive and behavioral development.

### Continuing Risks to Lead Exposure and Toxicity


Many of the most important risk factors are low income and socioeconomic status, houses containing lead-based paint, and poor nutrition. In a 1988 report to Congress, the Agency for Toxic Substances and Disease Registry (ATSDR) determined that about 10% of the 1 million or so black children that live in inner city areas of large U.S. cities and have an income below the poverty level have blood lead levels greater than 15  $\mu\text{g}/\text{dl}$  (15). Children who live in old houses with lead-containing paint and exposure to lead-containing dust are at increased risk for lead exposure and elevated blood lead levels. Lead was banned from indoor paint in the United States in 1978, so every house built before that time is likely to contain lead-based paint. This includes more than 50 million dwellings (3). It is estimated that at least 6 million children live in houses built prior to 1940—houses that constitute the greatest risk.

Less-than-ideal nutrition contributes to adverse consequences of lead exposure, particularly diets that are deficient in the essential minerals calcium, iron, and zinc, (16).

Studies with experimental animals fed diets low in calcium have established that calcium deficiency increases both tissue retention and toxicity of lead (17). Iron deficiency also increases tissue deposition and toxicity to lead (18). A high prevalence of iron deficiency occurs in infants and children because of the need to expand the body's iron pool during growth; it has been observed that the children with the highest environmental exposures to lead are also at the greatest risk to iron deficiency (19). The most significant health impact of iron deficiency is in young children who develop defects in attention span that lead to learning deficits (20). A longitudinal prospective study in Yugoslavia showed that both lead exposure and iron deficiency produce effects on neurobehavior and the hemopoietic system among pregnant women, infants, and children (21). Lead and zinc interactions are not as well defined as those between lead and calcium and between lead and iron. Experimental studies have shown that lead increases zinc excretion (22) and that zinc deficiency enhances lead absorption (23). There is also a close inverse relationship between blood lead levels and zinc-containing heme enzymes (24).

### Prenatal Exposures to Lead

Risk factors for prenatal exposure to lead involve maternal exposure and body burden of lead. There are both exogenous and endogenous factors contributing to maternal blood lead levels and *in utero* exposure to the fetus. Several studies have shown that there is no impairment to lead crossing the placenta and that maternal and fetal



**Figure 2.** Plot of time course of maternal blood lead (geometric mean and SE) during pregnancy for subjects with complete and incomplete data. Incomplete data are offset on the X-axis for clarity. Upward trend of complete subjects from 20 weeks to term is significant. Ranges of horizontal lines above the data indicate significant comparisons using least significant differences post-hoc test, with indicated probabilities. Conversion factor for blood lead is 10.0  $\mu\text{g}/\text{dl}$  = 0.48  $\mu\text{mol}/\text{l}$ . From Rothenberg et al. (26).

blood lead levels are similar (25). Today in the United States, environmental exposure to lead is largely from dietary sources for non-occupationally exposed women but, in areas of the world where leaded gasoline is still in use, airborne lead may be a significant source. Lead-glazed pottery from cottage industries may also be a potential source where such pottery is in use. Gastrointestinal absorption of lead may be increased during pregnancy along with increased calcium absorption.

The major endogenous source of lead is the skeletal system. Lead in bone reflects past exposure and may be independent of blood lead. The increase in the calcium requirement during pregnancy is accompanied by release of lead into blood which, in turn, results in fetal exposure to lead. An interesting demonstration of changes in blood lead levels during pregnancy was recently reported for 105 women living in Mexico City (26). The mean blood lead level for the group was 7.0  $\mu\text{g}/\text{dl}$  with a range of 1.3–35.5  $\mu\text{g}/\text{dl}$ , (Fig. 2). There is a significant decrease in blood lead level between 12 and 20 weeks (1.1  $\mu\text{g}/\text{dl}$ ). This has been explained by physiological factors such as increase in organ size and hemodilution. From week 20 to delivery, there is a linear upward trend.

Studies using stable isotope techniques have shown that about 45–75% of lead in blood is derived from bone lead in persons without excessive exposure to lead (27,28), and there is a further 32%–65% increase in the contribution of skeletal lead to blood lead during gestation (B.L. Gulson, personal communication). The wide variation is thought to be due to differences in the



ratio of bone lead to blood lead in specific individuals. It is expected that low dietary calcium will result in greater mobilization of bone calcium, and lead along with the calcium is contributed to supply needs of the fetus.

### Central Nervous System and Birth Outcomes from Prenatal Exposure to Lead

Assessment of cognitive development in relation to prenatal exposure to lead is difficult because of limitations in assessment methodology. Nevertheless, results of prospective epidemiological studies (29,30) have shown that children with high umbilical cord blood lead ( $>10 \mu\text{g}/\text{dl}$ ) have a slower sensorimotor or visual-motor development as measured by the Bayley Scales of Infant Development than children with lower blood lead levels at birth. Deficits in cognitive development with increased prenatal blood lead levels are likely to attenuate over time. Blood lead levels at 24 months of age have been found to be the most predictive of future cognitive development. Bellinger et al. (29) found that persistence of the cognitive deficit related to prenatal lead exposure is most often detected in those infants with either high postnatal exposure or less than optimal socioeconomic demographics.

Andrews et al. (31) reviewed 25 epidemiological studies to determine the relationship of prenatal lead exposure and birth outcomes. They concluded that prenatal lead exposure is unlikely to increase the risk of premature rupture of membranes but does appear to increase the risk of preterm delivery. It is unclear whether prenatal lead exposure decreases gestational age. Prenatal lead exposure appears to be related to reduced birth weight, but results

vary in relation to study design and degree of control for confounding factors.

### Exposure of Fetal Brain to Lead

Experimental studies have shown that the fetal blood-brain barrier is immature and uptake of lead by fetal rat brain during gestation is greater than after birth. There is a sixfold increase in total brain lead prenatally compared to a 3.5-fold increase during weaning and a twofold increase post-weaning with the same level of exposure (32). As animals mature, less lead enters the brain. Other experimental studies suggest that the immature endothelial cells forming the capillaries of the developing brain are less resistant to the effects of lead than capillaries from mature brains; these immature endothelial cells permit fluid and cations including lead to reach newly formed components of the brain, particularly astrocytes and neurons (33).

There are few studies of uptake of lead by the human brain during development. The largest number of lead measurements in fetal brain have been reported by Bartrop (34) in 21 human fetuses with maturities of 14–40 weeks, (Fig. 3). There was a direct relationship between brain weight and brain lead content, but there was no change in lead concentration. There was also a parallel increase in calcium; it has been suggested that movement of lead into the fetal brain, as well as other tissues, follows the movement of calcium. Studies in guinea pigs have shown that transfer of lead from mother to fetus is linearly related to umbilical cord blood flow rate (35).

### Mechanisms for Neurological Effects of Lead

Clinical and experimental studies have provided insights regarding the mechanisms whereby lead produces neurological effects on the developing brain as studied postnatally (36). Mechanisms for neurological effects as studied postnatally are likely to be a continuum of prenatal effects; however, it is not known whether there are unique prenatal neurological effects.


Effects of lead on the developing brain are both morphological and pharmacological.

- Neuropharmacological
  - Lead substitutes for calcium
  - Neurotransmitter Release
  - Protein kinase C
  - Na-Ca ATPase
  - Energy metabolism
- Neurodevelopmental (morphological)
  - Interference with adhesion molecules
  - Impaired programming of cell:cell connections
  - Miswiring of central nervous system (CNS).

One of the principal mechanisms by which the central nervous system develops during the early postnatal period is by overgrowth of neuronal processes, with subsequent "pruning" or deletion of synapses to adjust to the needs of its environment. For humans more brain development occurs after birth than in any other species. A highly significant morphological effect is the result of lead impairment of timed programming of cell:cell connections resulting in modification of neuronal circuitry (37). This effect has been shown to be the result of decreased sialic acid production by neural cells, which produces a failure in synaptic structuring (38). This disruption of the toning or matching process of neuronal connections or "miswiring" may produce functional effects. It has also been found that lead induces precocious differentiation of the glia upon which cells migrate to their eventual positions during structuring of the brain, further enhancing the likelihood of alterations in normal development (39).

Lead functions pharmacologically by interfering with synaptic mechanisms for neurotransmitter release. It has been suggested that lead can substitute for calcium and possibly zinc in ion-dependent events at the synapse and is responsible for the observed impairment by lead of various neurotransmitter systems (cholinergic, GABergic, and dopaminergic) (36,40). The neonatal period has been found to be most sensitive to inhibitory action of lead on *N*-methyl-D-aspartate receptor-ion function channels (41). *In vitro* studies on brain capillaries have shown that micromolar concentrations of lead activate protein kinase C, a second messenger in the regulation of cellular metabolism (42). Lead may also replace calcium in calmodulin-dependent reactions, inhibit membrane-bound  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase, and interfere with mitochondrial release of calcium with impairment of energy metabolism (43). These effects are potentially reversible if lead can be removed from active sites. Although removal from lead exposure and chelation therapy lower blood lead levels, there is little or no information regarding effects from removal of lead from these sensitive molecular sites (44).

In spite of the demonstrated morphological and pharmacological effects of lead in the maturing brain, it is difficult to demonstrate a functional or clinical effect in the child under 2 years of age because of the immaturity of the brain and limited number of functions that can be tested at that age. There is no consistent lead neuropsychological syndrome or behavioral signature that can be identified clinically (45). In some studies, lead is most closely



**Figure 3.** Relationship between brain weight and brain lead content in 21 human fetuses with maturities of 14–40 weeks. From Bartrop (34).

related to loss in verbal skills; in others, it is nonverbal skills. Studies of cohorts of children with postnatal exposure to lead determined that the perceptual-motor subsets of the McCarthy Scales of Children's Abilities were particularly sensitive to lead exposure (46). Lead related losses in visual motor integration, poorer performance on tasks of serial learning, and increased errors in tasks requiring perseverance in experimental animal models are generally consistent with clinical findings and are thought to pinpoint the hippocampus as the primary target for neurobehavioral effects (47–50). Also, the hippocampus is known to accumulate lead in humans (51) and rat pups exposed to lead (52). However, Lilienthal et al. (53) found that the behavioral profile in rats with chemically induced lesions of the hippocampus differed from patterns obtained in rats with prenatal exposure to lead and that chemical lesions of the amygdala show a greater similarity with lead-induced effects, suggesting that the hippocampus is not the sole target for lead toxicity.

## Summary

Advances in the understanding of the toxic effects of lead is an outstanding example of how knowledge learned from research can impact public health. Reduction of lead in gasoline, elimination of lead solder from canned food, removal of lead from paint, and abatement of housing containing lead-based paint are important contributions to reduction of lead exposure. Factors that enhance risk to lead exposure, particularly during fetal life, are low socioeconomic status, old housing with lead-containing paint, and less than ideal nutrition, particularly low dietary intake of calcium, iron, and zinc. Prenatal exposure may result from endogenous sources such as lead in the maternal skeletal system or maternal exposures from diet and the environment.

The developing nervous system is particularly sensitive to the toxic effects of lead, and experimental studies have shown that a large number of the effects in the nervous system are due to interference by lead with biochemical functions dependent on calcium ions and impairment of neuronal connections dependent on dendritic pruning.

In spite of past efforts, there are many important avenues for further research, particularly with regard to prenatal exposures and neurological effects. Ongoing studies using stable-isotope techniques should provide new information regarding maternal–fetal transfer of lead. There is need for a better understanding of the mechanisms for the toxic effect of lead on

the nervous system and also to determine whether effects from postnatal exposure are a continuum of prenatal lead exposure or whether prenatal exposure to lead produces unique effects.


## REFERENCES

- Goyer RA. Lead toxicity: current concerns. *Environ Health Perspect* 100:177–187 (1993).
- U.S. EPA. Air quality criteria for lead, vols I–IV. EPA-600/8-83/028f. Washington:U.S. Environmental Protection Agency, 1986.
- National Research Council. Measuring lead exposure in infants, children and other sensitive populations. Washington:National Academy Press, 1993.
- International Program for Chemical Safety. Inorganic lead. Environmental Criteria Document 165. Geneva:World Health Organization, 1995.
- CDC. Preventing lead poisoning in young children: a statement by the centers for disease control, Atlanta, GA:Centers for Disease Control, 1991.
- Health Canada. Update of evidence for low-level effects of lead and blood lead intervention levels and strategies—final report of working group, Ottawa, Canada:Environmental Health Directorate, 1994.
- Proceedings of the International meeting on non-occupational exposure to lead, 5–9 October 1992. Melbourne, Australia (Edwards-Bert P, Bently K, Callan P, Baghurst P, eds). Adelaide:South Australian Health Commission and Commonwealth Department of Health Housing and Local Government Services, 1993.
- Cory-Slechta DA. Bridging human and experimental animal studies of lead neurotoxicity: moving beyond IQ. *Neurotoxicol Teratol* 17:219–221 (1995).
- CDC. Strategic plan for the elimination of childhood lead poisoning. Washington:U.S. Department of Health and Human Services, 1991.
- HUD. Comprehensive and workable plan for the abatement of lead-based paint in privately owned housing: report to Congress. Washington:U.S. Department of Housing and Urban Development, 1990.
- EPA. Strategy of reducing lead poisoning, Washington:U.S. Environmental Protection Agency, 1991.
- Bolger PM, Carrington CD, Capar SG, Adams MA. Reductions in dietary lead in the United States. *Chem Spec Bioavail* 314:31–36 (1991).
- Guthrie AM, McNulty M. Making the most of Medicaid: state progress in childhood lead poisoning prevention. Washington:Alliance to End Childhood Lead Poisoning, 1993.
- Pirkle JL, Brody DJ, Gunter EW, Kramer RA, Paschal DC, Flegal KM, Matte TD. The decline in blood lead levels in the United States. *JAMA* 272:284–291 (1994).
- ATSDR. The nature and extent of lead poisoning in the United States. A report to Congress. Atlanta, GA:Agency for Toxic Substances and Disease Registry, 1988.
- Goyer RA. Nutrition and metal toxicity. *Am J Clin Nutr* 61 (suppl):646S–650S (1995).
- Mahaffey KR. Factors modifying susceptibility to lead toxicity. In: *Dietary and environmental lead: human health effects* (Mahaffey KR, ed). New York:Elsevier, 1985:373–420.
- Six KM, Goyer RA. The influence of iron deficiency on tissue content and toxicity of ingested lead in the rat. *J Lab Clin Med* 79:128–136 (1972).
- Mahaffey KR, Annett JL. Association of erythrocyte protoporphyrin with blood lead level and iron status in the Second National Health and Nutrition Examination Survey, 1976–1980. *Environ Res* 41:327–338 (1986).
- Pollitt EP, Hathirat NJ, Kutchabhaki L, Missell L, Valyasevi A. Iron deficiency and educational achievement in Thailand. *A J Clin Nutr* 50(suppl 3):687–697 (1989).
- Graziano JH, Popovac D, Factor-Litvak P, Shrout P, Kline J, Murphy MJ, Zhao Y, Mehmeri A, Ahmed X, Rajovic B. Determinants of elevated blood lead during pregnancy in a population surrounding a lead smelter in Yugoslavia. *Environ Health Perspect* 89:95–100 (1990).
- Victory W, Thomas D, Shoeps P, Vander AJ. Lead increases urinary zinc excretion in rats. *Biol Trace Elem Res* 4:211–219 (1982).
- Petering HG. Some observations on the interaction of zinc, copper, and iron metabolism in lead and cadmium toxicity. *Environ Health Perspect* 25:141–145 (1978).
- Dutkiewicz B, Dutkiewicz T, Milkowska G. The effect of mixed exposures to lead and zinc on ALA level in urine. *Int Arch Occup Environ Health* 42:341–348 (1979).
- Goyer RA. Transplacental transport of lead. *Environ Health Perspect* 89:101–108 (1990).
- Rothenberg SJ, Karchmer S, Schnaas L, Perroni E, Zea F, Alba JF. Changes in serial blood lead levels during pregnancy. *Environ Health Perspect* 102:876–880 (1995).
- Gulson BL, Mahaffey KR, Mizon KJ, Korsch MJ, Cameron MA, Vimpani G. Contribution of tissue lead to blood lead in adult female subjects based on stable lead isotope methods. *J Lab Clin Med* 125:703–712 (1995).
- Smith DR, Osterloh JD, Flegal AR. Use of endogenous, stable lead isotopes to determine release of lead from the skeleton. *Environ Health Perspect* 104:60–66 (1996).
- Bellinger D, Leviton A, Waternaux C, Needleman H, Rabinowitz M. Longitudinal analyses of prenatal and postnatal lead exposure and early cognitive development. *N Engl J Med* 316:1037–1043 (1987).
- McMichael AJ, Baghurst PA, Wigg NR, Vimpani GV, Robertson EF, Roberts RJ. Port Pirie cohort study: environmental exposure to lead and children's abilities at the age of four years. *N Engl J Med* 319:468–475 (1988).
- Andrews KW, Savitz DA, Hertz-Picciotto I. Prenatal lead exposure in relation to gestational age and birth weight: a review of epidemiological studies. *Am J Ind Med* 26:13–32 (1994).
- Rossouw J, Offermeier J, Van Rooyen JM. Apparent central neurotransmitter receptor changes induced by low-level lead exposure during different developmental phases in the rat. *Toxicol Appl Pharmacol* 91:132–139 (1987).
- Toews AD, Kolber A, Hayward J, Krigman MR, Morrell P. Experimental lead encephalopathy in the suckling rat: concentration of lead in cellular fractions enriched in brain capillaries. *Brain Res* 147:131–138 (1978).
- Bartrop D. Transfer of lead to the human fetus. In: *Mineral metabolism in pediatrics* (Bartrop D, Burland W, eds). Oxford: Blackwell Scientific

- Publications, 1969;135-151.
35. Kelman BJ, Walter BK. Transplacental movements of inorganic lead from mother to fetus. *Proc Soc Exp Biol Med* 163:278-282 (1980).
  36. Silbergeld EK. Mechanisms of lead neurotoxicity, or looking beyond the lampost. *FASEB J* 6:3201-3206 (1992).
  37. Bull RJ, McCauley PT, Taylor DH, Croften KM. The effects of lead on the developing nervous system of the rat. *Neurotoxicology* 4:1-18 (1983).
  38. Cookman GR, Hemmens SE, Keane GJ, King WB, Regan CM. Chronic low-level lead exposure precociously induces rat glial development *in vitro* and *in vivo*. *Neurosci Lett* 86:33-37 (1988).
  39. Cookman GR, King W, Regan CM. Chronic low-level lead exposure impairs embryonic to adult conversion of the neural adhesion molecule. *J Neurochem* 49:399-403 (1987).
  40. Verity MA. Nervous system. In: *Metal toxicology* (Goyer RA, Klaassen CD, Waalkes MP, eds). San Diego, CA:Academic Press, 1995;199-226.
  41. Guilarte TR, Miceli RC, Altman L, Weinsberg F, Winneke G, Wiegand H. Chronic prenatal and postnatal Pb+2 exposure increases [<sup>3</sup>H]MK801 binding sites in adult rat forebrain. *Eur J Pharmacol* 248:273-275 (1993).
  42. Markovac J, Goldstein GW. Lead activates protein kinase C in immature rat brain microvesicles. *Toxicol Appl Pharmacol* 96:14-23 (1988).
  43. Simons TJB. Cellular interactions between lead and calcium. *Br Med Bull* 42:431-434 (1986).
  44. Goyer RA, Cherian MG, Jones MM, Reigart JR. Role of chelating agents for prevention, intervention, and treatment of exposures to toxic metals. *Environ Health Perspect* 103:1048-1053 (1995).
  45. Bellinger DC. Interpreting the literature on lead and child development: the neglected role of the "experimental system." *Neurotoxicol Teratol* 17:201-212 (1995).
  46. Wasserman GA, Graziano JH, Factor-Litvak P, Popovac D, Morina N, Musabegovic A, Vrenezi N, Capuni-Paracka S, Lekic V, Preteni-Redjepi E. Consequences of lead exposure and iron supplementation on childhood development at age 4 years. *Neurotoxicol Teratol* 16:233-240 (1994).
  47. Baghurst PA, McMichael AJ, Tong S, Wigg NR, Vimpani GV, Robertson EF. Exposure to environmental lead and visual-motor integration at age 7 years: the Port Pirie cohort study. *Epidemiology* 6:104-109 (1995).
  48. Davis JM, Otto DA, Weil DE, Grant LD. The comparative developmental neurotoxicity of lead in humans and animals. *Neurotoxicol Teratol* 12:215-229 (1990).
  49. Rice DC. Behavioral effects of lead in monkeys tested during infancy and adulthood. *Neurotox Teratol* 14:235-245 (1992).
  50. Alfano DP, Petit TL. Behavioral effects of postnatal lead exposure: possible relationship to hippocampal dysfunction. *Behav Neural Biol* 32:319-333 (1981).
  51. Grandjean P. Regional distribution of lead in human brains. *Toxicol Lett* 2:65-69 (1978).
  52. Lafauconnier JM, Bernard G, Mellerio F, Seville A, Cesarini E. Lead distribution in the nervous system of 8-month old rats intoxicated since birth with lead. *Exp* 39:1030-1031 (1989).
  53. Lillenthal H, Munoz C, Winneke G, Lenaerts C, Hennekes R. Neurobehavioral and electrophysiological effects of lead in rats and monkeys. In: *Lead exposure and child development* (Smith MA, Grant LD, Sors AI, eds). Boston:Kluwer Academic Publishers, 1989;414-426.

# FIRST INTERNATIONAL CONFERENCE ON EMERGING ZOOSES

Holiday Inn Crowne Plaza  
Jerusalem, Israel  
November 24-28, 1996



Co-Sponsored by the Centers for  
Disease Control

For further information:  
Emerging Zoonoses  
PO Box 29041  
Tel Aviv 61290  
Israel  
972 3 5175149/50  
Fax: 972 3 517 5155